

The Radionactive Universe

Edward W. Kolb

Theoretical Astrophysics Group, Fermi National Accelerator Laboratory, Box 500, Batavia, Illinois 60510, and Department of Astronomy and Astrophysics, Enrico Fermi Institute, The University of Chicago, Chicago, Illinois 60637
E-mail: rocky@fnal.gov

Géraldine Servant

High Energy Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, and Enrico Fermi Institute, The University of Chicago, Chicago, Illinois 60637
E-mail: servant@theory.uchicago.edu

Tim M. P. Tait

Fermi National Accelerator Laboratory, Box 500, Batavia, Illinois 60510
E-mail: tait@fnal.gov

ABSTRACT: In a large class of extra-dimensional models, a scalar degree of freedom known as the *radion* is long-lived, or even stable, on cosmological scales. In this paper we investigate the impact of radionactivity on the evolution of the universe. We demonstrate that whether the radion overcloses the universe, constitutes the dark matter, is the inflaton, the curvaton, or does not play any role in cosmology, depends crucially on the ratio between the energy densities stored in the radion and in the inflaton at the time of inflation. We discuss the general difficulties reconciling models with low compactification scale (*i.e.*, TeV scale) with the simple picture of inflation.

KEYWORDS: Cosmology of Theories beyond the SM, Physics of the Early Universe.

Contents

1. Introduction	2
2. Radion Properties	5
2.1 Radion Mass	7
2.2 Radion Couplings	7
2.3 Radion Lifetime	8
2.4 Radion as the Inflaton	8
3. Radion Mixing and Induced Inflationary Potential	10
4. Evolution of the Radion Field	13
4.1 Evolution During Inflation	13
4.2 Validity of 4D effective field theory	17
5. Radion as Dark Matter	17
5.1 Coherent Production	17
5.2 Thermal Production	19
6. Radions and Cosmological Perturbations	21
6.1 Radion Isocurvature Perturbations	21
6.2 Radion as the Curvaton	22
6.3 Radion as the Modulon	23
7. Late Decay of Radions and KK Gravitons	25
7.1 Late Decay of Radions	25
7.2 Late Decay of KK gravitons	26
7.3 Diffuse Gamma-ray Signal	28
8. Conclusions	28
8.1 TeV Extra Dimensions	30

1. Introduction

Scalar fields are believed to play important roles in both particle physics and cosmology. The crucial aspect of scalar fields is that they determine the properties of the vacuum without relinquishing the key property: invariance under spacetime symmetries. In spite of these very desirable properties, scalar fields fit rather uneasily within quantum field theories. Scalar field masses are extremely sensitive to quantum corrections. They exhibit quadratic divergences, which suggests that their natural mass scale should be very large, and implies that their detailed properties should be controlled by ultraviolet (UV) physics beyond our ability to infer from current experiments. The large masses natural for scalar fields tend to spoil their potential utility in both particle physics and cosmology. Ideally, we should be led to consider theories with some mechanism to protect the scalar mass from large corrections.

In this article we explore one generic feature of theories with extra spatial dimensions. Such theories contain a scalar field, the radion, which plays the role of the geometrical modulus that determines the size of the compactified dimensions. From the higher dimensional perspective, the radion is part of the enlarged graviton tensor, and at distances smaller than the size of the extra dimension it is protected from receiving a large mass by the same general covariance which protects the graviton mass. Thus, unlike generic scalar fields, it can be naturally light. Light scalar degrees of freedom are often invoked in cosmology: inflatons, curvatons, modulons (fields controlling coupling strengths), dark matter, *etc.* It is interesting to explore the possibility that the radion may be one of the much-needed light scalar degrees of freedom, and to see if it can play an interesting role in cosmology.

Extra-dimensional models with a TeV compactification scale [1] have received revived interest lately. There are many ways in which extra dimensions can attempt to explain the outstanding mysteries in particle physics [2, 3, 4]. For example, there is the prospect of dark matter consisting of the first Kaluza–Klein (KK) mode of the hypercharge gauge boson (B^1) [5] in a sub-class of these models referred to as *universal extra dimensions* (UED) [6]. However, the cosmology of models with TeV size extra dimensions remains largely unexplored. Our concern in this paper is to investigate, for all possible compactification scales, constraints from radion cosmology on these models, and to a larger extent, on any models with flat extra dimensions. Different recent attempts discussing the role of the radion during inflation can be found in [7].

The destiny of a radionactive universe depends on two parameters, the compactification scale M_c (naturally L^{-1} , the physical size of the extra dimensions) and the ratio $r = M_c/M_I$ where $M_I = V_I^{1/4}$ is the scale of the inflaton potential. One question to ask concerns the epoch at which the extra dimensions were stabilized compared to the epoch of inflation. When did compactification take place relative to inflation? A first possibility is that inflation took place before compactification. This would require a post-inflationary exponential contraction of extra dimensions, and seems counterintuitive. Further, it would require that the inflation scale M_I is much larger than the compactification scale M_c . As is well known, the non-renormalizability of extra-dimensional theories implies that they have a low cut-off scale Λ ,

beyond which the theory is strongly coupled and loses predictivity.¹

The second possibility stabilizes the extra dimensions, and then inflates the four large ones much later. In this case, geometrical moduli such as the radion are active during inflation. These are scalar fields with Planck-suppressed couplings, and may contribute to the inflaton potential. In fact, they may be responsible for triggering inflation, for all possible compactification scales [9]. Even if they do not have the properties required to trigger inflation, they may still play a crucial role in the generation of cosmological perturbations and/or make up the dark matter. As we shall see, these scalar fields are typically light, with masses protected by higher dimensional general covariance. It is well known that when the mass is smaller than the Hubble rate during inflation, the field is frozen during inflation. After inflation, the energy density stored in these fields can easily overclose the universe. This is the essence of the moduli problem. Another well known fact is that during inflation, scalar fields receive corrections to their mass of order H from non renormalizable operators of the type $V_I \phi^2/m_{Pl}^2$ (where ϕ is the modulus in question). Scalars with masses larger than approximately H are damped out to the minimum of their potential during inflation. However, their energy is not necessarily inflated away because the position of the minimum is shifted by the same type of operators which are responsible for shifting the masses. As a result, energy stored in the radion fields typically survives inflation and can have dramatic effects on post-inflationary cosmology. We can actually predict m^2/H^2 during inflation as a function of r . We show that if $r \lesssim 1$, models with a compactification scale $M_c \lesssim 10^9$ GeV can easily be ruled out (at least within a given inflationary paradigm) because of the overclosure of the universe by radion oscillations. This means that inflation must take place at a scale as low as or lower than the compactification scale.

Inflation taking place after compactification appears problematic in models with a low compactification scale, *i.e.*, $M_c \sim \text{TeV}$. This implies that the highest energy scale of the universe when inflation starts is $M_I \lesssim M_c$. From the COBE constraint $\delta_H \sim 10^{-5}$ [10] and by identifying the energy density fluctuations with primordial quantum fluctuations of the inflaton, one obtains:

$$\left(\frac{V_I}{\epsilon}\right)^{1/4} \sim 10^{16} \text{ GeV}, \quad (1.1)$$

where $\epsilon = m_{Pl}^2(V_I'/V_I)^2/2$ is the slow-roll parameter. This data disfavors models with low-scale inflation. Equation (1.1) tells us that if the primordial density perturbations originate from the inflaton, inflation at the TeV scale requires a slow-roll parameter of the order of 10^{-48} , a tiny value which is difficult to motivate naturally. However, it has recently been hypothesized that structures in the universe may not necessarily follow from primordial quantum fluctuations in the inflaton, but could instead be due to another scalar field, referred to as the *curvaton* [11]. The curvaton scenario has the advantage that the constraints on the inflaton potential are relaxed, though at the cost of introducing another scalar field whose mass must

¹For theories in which gauge fields propagate in the bulk, naïve analysis suggests $\Lambda \sim 100M_c$ (see however [8] which suggests $\Lambda \sim M_c$) if there are 5 dimensions, and Λ is closer to M_c for theories with more dimensions. The scale at which gravitational interactions become strong is typically about $M_5^3 \sim M_c m_{Pl}^2 \gg \Lambda^3$ in 5d.

somehow be protected. Furthermore, the constraints are not relaxed to the point that TeV inflation becomes natural. Indeed, the COBE constraint in the curvaton scenario translates into

$$\frac{f_D H_*}{\phi_*} \approx 10^{-4}, \quad (1.2)$$

where H_* and ϕ_* are the Hubble constant and the amplitude of the curvaton field respectively at the time of horizon exit. Here f_D is the fraction of the total energy density contained in the curvaton when it decays and should be of order unity for this mechanism to work. To satisfy Eq. (1.2), an inflation scale at a TeV would require the small value $\phi_* \sim 10^{-9}$ GeV $\ll M_I$ for the curvaton amplitude, which is hard to reconcile with the possibility that the curvaton dominates the total energy density at the time of its decay. An even more recent proposal [12] suggests that cosmological perturbations could be due to primordial fluctuations in the field (which we will refer to as a *modulon*) whose VEV gives the coupling between the inflaton and the Standard Model fields and plays a crucial role in the reheating process. In this work we will investigate whether the radion could ever play the role of a curvaton or a modulon.

A third alternative has compactification and inflation taking place at comparable scales. We assume that the relevant history of our universe starts at a time when all dimensions have comparable sizes. Inflation starts in all $4 + n$ dimensions (with a typical scale around or a few orders of magnitude below m_{Pl}). Later on, the extra dimensions stop inflating, for example when M_c is at the TeV scale, and the rest of the 4 dimensions continue to inflate. One assumes there are some extra fields living in the extra dimensions responsible for the stabilization dynamics and allowing the extra dimensions to undergo only 15 – 20 e-folds of inflation while allowing the other dimensions to continue to inflate. In this case, the scale in the inflaton potential is still much higher than the compactification scale, $M_I \gg M_c$, and there would be no need for a huge fine-tuning in the epsilon parameter or in ϕ_* to get the right COBE normalization. As we will see in section 5, there is a serious moduli problem associated with this scenario. Indeed, when the ratio between the compactification scale and the inflaton scale is much smaller than one, we are in the regime where there is no damping at all of the radion field during inflation, and it will overclose the universe. Avoiding the overclosure problem typically requires $M_I \ll M_c$, which on the other hand disfavors low compactification scales from the COBE constraint, unless there is a new mechanism to generate density perturbations. Obviously, there are interesting cosmological challenges associated with TeV compactification scales. In this work, in addition to examining the difficulties associated with low compactification scales, we will be able to constrain even larger compactification scales, up to 10^{12} GeV. This leads us to study of more traditional compactifications models, *i.e.*, associated with smaller extra dimensions.

In summary, as soon as compactification took place before inflation, we are presented with additional scalar fields during inflation, whose masses are protected from large corrections, and thus they typically survive inflation. Note that as the number of extra dimensions increases, the number of geometric moduli also increases. We concentrate on the volume-modulus, *i.e.*, we focus on the 5d case for illustrative purposes. In principle, one may also consider more

extra dimensions, and those moduli which describe the shape of the compact manifold.

Section 2 reviews the definition and properties of the radion field and contains a few remarks about the possibility that the radion is itself the inflaton. Section 3 stresses the consequences of the generic form of the radion coupling to any form of energy density and how the radion can be naturally induced to be an inflaton. In section 4 we evaluate the energy density stored in the radion at the end of inflation and discuss its evolution when it begins to oscillate. For small enough M_c , the radion is stable on the time-scale of the age of the universe, and could either account for the dark matter, or overclose the universe. This is explored in section 5. In section 6, after studying the radion isocurvature perturbations, we discuss whether the radion can be a viable curvaton or modulon candidate, and can account for the observed density perturbations. Section 7 studies the constraints associated to the late decay (after BBN) of radions and KK gravitons. In section 8 we summarize the challenges associated with TeV compactification scales and conclude.

2. Radion Properties

In this section, we briefly review the crucial properties of the radion: its mass, and its coupling to bulk fields. When reducing the curvature term of the $(\hat{d} = d + n)$ -dimensional action, we obtain terms of the form

$$\begin{aligned} -\frac{1}{\hat{\kappa}^2} \int d^{\hat{d}}x \sqrt{g^{(\hat{d})}} \mathcal{R}^{(\hat{d})} &= \int d^d x \sqrt{g^{(d)}} \sqrt{g^{(n)}} e^{\frac{d-2}{2}\Sigma} \left[-\frac{1}{\kappa^2} \mathcal{R}^{(d)} + \frac{1}{4} e^{-\Sigma} g_{ij} F_{\mu\nu}^i F^{j\mu\nu} \right. \\ &\quad - \frac{(d-1)(d-2)}{4\kappa^2} (\partial_\mu \Sigma)(\partial^\mu \Sigma) - \frac{d-1}{2\kappa^2} g^{ij} (\partial_\mu g_{ij})(\partial^\mu \Sigma) \\ &\quad \left. + \frac{1}{4\kappa^2} g^{ij} (\partial_\mu g_{jk}) g^{kl} (\partial^\mu g_{li}) - \frac{1}{4\kappa^2} g^{ij} (\partial_\mu g_{ij}) g^{kl} (\partial^\mu g_{kl}) + \dots \right] \end{aligned} \quad (2.1)$$

where the general metric is given by

$$ds^2 = (e^\Sigma g_{\mu\nu} + 4g_{ij} \kappa^2 A_\mu^i A_\nu^j) dx^\mu dx^\nu + g_{ij} (2\kappa A_\mu^i dx^\mu dy^j + 2\kappa A_\nu^j dx_\nu dy^i) + g_{ij} dy^i dy^j \quad . \quad (2.2)$$

All fields are functions of both extra-dimensional coordinates y_i and four-dimensional coordinates x^μ . However, the zero-mode fields are independent of the y_i , and thus we may drop this dependence when describing the low-energy (zero-mode) physics. In five dimensions, the KK tower of the radion is eaten by the massive KK gravitons leaving only the radion zero mode; in more extra dimensions, some of the radion KK modes remain as well. $\hat{\kappa}$ is Newton's constant in $\hat{d}$ dimensions and is related to Newton's constant in d (flat) dimensions ($\kappa = m_{Pl}^{-2}$ where m_{Pl} is the reduced 4-d Planck mass) by $\kappa = \hat{\kappa}/\sqrt{V_n}$ where V_n is the volume of the compact space. Here $g^{(d)}$ and $g^{(n)}$ are the determinants of the metrics $g_{\mu\nu}$ and g_{ij} . In the right-hand side, μ and ν indices go up and down with the $g_{\mu\nu}$ metric and its inverse. To obtain a curvature term in d dimensions which is canonically normalized, we make the following rescaling:

$$\sqrt{g^{(n)}} e^{\frac{d-2}{2}\Sigma} = 1, \quad i.e., \quad e^\Sigma = 1/\sqrt{g^{(n)}} \quad . \quad (2.3)$$

In this case, Eq. (2.1) simplifies to

$$-\frac{1}{\hat{\kappa}^2} \int d^{\hat{d}}x \sqrt{g^{(\hat{d})}} \mathcal{R}^{(\hat{d})} = \int d^d x \sqrt{g^{(d)}} \left[-\frac{1}{\kappa^2} \mathcal{R}^{(d)} + \frac{1}{4} g^{(n)\frac{1}{d-2}} g_{ij} F_{\mu\nu}^i F^{j\mu\nu} \right. \\ \left. + \frac{1}{4\kappa^2} g^{ij} (\partial_\mu g_{jk}) g^{kl} (\partial^\mu g_{li}) + \frac{1}{4(d-2)\kappa^2} g^{ij} (\partial_\mu g_{ij}) g^{kl} (\partial^\mu g_{kl}) + \dots \right]. \quad (2.4)$$

We now apply these results to the 5-dimensional case, S^1/Z_2 with bulk fields. The 5d action can be expressed as

$$\mathcal{L}_5 = \sqrt{g} \left\{ M_5^3 R_5 - \frac{1}{4g_5^2} \text{Tr}[F^{MN} F_{MN}] - M_c^5 \tilde{V}_\phi(\tilde{\phi}) - \frac{1}{2} M_\psi^3 \partial_M \tilde{\psi} \partial^M \tilde{\psi} - M_\psi^5 \tilde{V}_\psi(\tilde{\psi}) \right\}, \quad (2.5)$$

where g , M_5 , and R_5 refer to the 5d gravitational quantities, F^{MN} is the field strength of a bulk gauge field with 5d coupling g_5 , $\tilde{\psi}$ represents a minimally coupled real bulk scalar with potential $\tilde{V}_\psi$, and $\tilde{V}_\phi$ is the radion potential. For simplicity, we have neglected the possibility that $\tilde{\psi}$ mixes with the Ricci scalar; we will return to this later.

A convenient parametrization for the 5d metric can be written in the form [13]:

$$ds^2 = \left(e^{-1/3\tilde{\phi}} g_{\mu\nu} + e^{2/3\tilde{\phi}} \tilde{A}_\mu \tilde{A}_\nu \right) dx^\mu dx^\nu + 2e^{2/3\tilde{\phi}} \tilde{A}_\mu dx^\mu dy + e^{2/3\tilde{\phi}} dy^2, \quad (2.6)$$

where $\tilde{A}_\mu = 2\kappa A_\mu$ is the gravi-photon field, which is odd under the orbifold and thus has no zero mode, $g_{\mu\nu}$ is the tensor graviton field, and $\tilde{\phi}$ is the (dimensionless) gravi-scalar (radion). This form is a convenient starting point because it decouples the kinetic terms of the radion and the transverse tensor modes and produces a radion kinetic term of the canonical form.

To determine the effective action at scales less than of order $1/L$, we replace the 5d fields with their zero-mode components, and integrate out the extra dimension. In doing so, it is convenient to introduce a reference length L . Note that the physical size of extra dimensions is $L_{phys} = e^{\tilde{\phi}/3} L$. The effective action for the zero modes is,

$$\mathcal{L}_4 = \sqrt{g} \left[M_5^3 L R_4 - \frac{M_5^3 L}{6} \partial_\mu \tilde{\phi} \partial^\mu \tilde{\phi} - \frac{L}{4g_5^2} e^{\tilde{\phi}/3} F_{\mu\nu} F^{\mu\nu} - \frac{M_\psi^3 L}{2} \partial_\mu \tilde{\psi} \partial^\mu \tilde{\psi} \right. \\ \left. - M_\psi^5 L e^{-\tilde{\phi}/3} \tilde{V}_\psi(\tilde{\psi}) - M_c^5 L \tilde{V}_\phi(\tilde{\phi}) \right], \quad (2.7)$$

where now all quantities refer to the 4d (zero-mode) fields. From here it is simple to rescale fields to canonical normalization and define four-dimensional scalar potentials,

$$\phi = \sqrt{M_5^3 L/3} \tilde{\phi}, \quad \psi = \sqrt{M_\psi^3 L} \tilde{\psi}, \\ V_\phi(\phi) = L M_c^5 \tilde{V}_\phi(\tilde{\phi}), \quad V_\psi(\psi) = L M_\psi^5 \tilde{V}_\psi(\tilde{\psi}). \quad (2.8)$$

In order for the low energy physics to appear four dimensional, the radion potential must provide a stable VEV for ϕ . We can adjust the reference quantity L such that this happens for $\langle \phi \rangle = 0$. Clearly, unless there is fine-tuning in the radion potential, this will occur for

$L \sim M_c^{-1}$. For this choice, L is in fact the physical size of the extra dimension, and we can identify $M_5^3 L = m_{Pl}^2$ and $g_5^2/L = g_4^2$. The radion couplings are thus,

$$-\frac{1}{4}e^{\phi/\sqrt{3}m_{Pl}}F_{\mu\nu}F^{\mu\nu} - e^{-\phi/\sqrt{3}m_{Pl}}V_\psi(\psi), \quad (2.9)$$

where we have rescaled the kinetic terms for the bulk fields to canonical normalization as in Eq. (2.8).

2.1 Radion Mass

In the absence of contributions from V_ψ , which we consider in the next section, the radion mass may be determined by expanding the potential $\tilde{V}_\phi$ appearing in Eq. (2.7) to order $\tilde{\phi}^2$ about its minimum,

$$-M_c^5 L \frac{1}{2} \tilde{V}_\phi''(\tilde{\phi}) \tilde{\phi}^2 + \dots = -M_c^5 L \frac{1}{2} \tilde{V}_\phi''(\tilde{\phi}) \frac{3}{m_{Pl}^2} \phi^2 + \dots \quad (2.10)$$

where primes refer to functional derivatives with respect to $\tilde{\phi}$. Without fine tuning, one expects $L \sim M_c^{-1}$, and all derivatives of $\tilde{V}_\phi$ to be order unity. Thus, the radion mass is naturally

$$m_r \sim \sqrt{3} \frac{M_c^2}{m_{Pl}}. \quad (2.11)$$

Note that loop corrections will not destabilize the radion mass, as they are of order M_c^2/m_{Pl} . Indeed they are proportional to the radion coupling and they must involve M_c since in the limit of decompactification the radion is part of the higher dimensional graviton, and thus is massless. The fact that the radion is typically light and its mass protected was stressed in [14] where it was also argued that for $M_c \sim \text{TeV}$, the radion would be responsible for detectable deviations to Newtonian gravity in present or future experiments. Thus, there is a lower bound on M_c from short-range gravitational experiments [14, 15, 16]. A radion with a mass of 10^{-3} eV would induce modification of Newton's law at distances of order $100 \mu\text{m}$ (at the edge of distances probed by gravitational experiments). No deviation has been measured, therefore

$$m_r \gtrsim 10^{-3} \text{ eV} \rightarrow M_c \gtrsim 0.8 \text{ TeV}, \quad (2.12)$$

assuming the gluons are bulk fields. The limits for brane gluons are weaker as a result of a weaker coupling between brane nucleons and the radion (see below). In these cases, direct collider limits will provide a lower bound on M_c of order 1 TeV, the precise value depending on the extra-dimensional framework [17].

2.2 Radion Couplings

Radion couplings to bulk fields may be simply read from Eq. (2.9), and are characterized by the four dimensional Planck scale, m_{Pl} . For much of the parameter space, $M_c \lesssim 10^8 \text{ GeV}$, the mass of the radion is less than 1 MeV, and decays proceed only into photons (and zero-mode

gravitons). These results formally apply only to flat extra dimensions and bulk SM fields. Two other possibilities are brane world models [18] in flat extra dimensions, or warped (RS) extra dimensions [19]. In the RS case, the radion mass and couplings are both characterized by the TeV scale [20, 21]. This large mass will ruin most of the cosmological applications of the radion we are considering, and thus we will not consider the warped case any further (see however, Ref [22]).

For flat extra dimensions with SM fields on a brane, the radion couplings are substantially different because brane fields do not contribute to the fifth components of the stress-energy tensor, and thus couplings to radions are induced only through the mixing of g_{55} with longitudinal gravitons [23]. The net result is that the radion couples only to the trace of the brane stress-energy tensor. For gauge fields, this occurs at tree-level proportionally to the gauge-field mass, and at one loop proportionally to the β function. The brane photon thus couples to the radion more weakly than a bulk photon would, with a relative coupling that is roughly $16\pi^2$ smaller. We parameterize the radion coupling to SM fields by c/m_{Pl} , with $c = 1$ for the bulk SM, and $c \sim 10^{-3}$ for the brane SM case. However, zero-mode gravitons, as bulk fields, will always couple with full strength to the radion, and thus the brane photon case does not substantially affect the radion lifetime, but instead suppresses the branching ratio for decay into photons. We will focus on the bulk photon case below, but occasionally comment on the brane-world scenario where appropriate.

2.3 Radion Lifetime

The decay into two bulk photons implies a radion decay width

$$\Gamma = \tau^{-1} \simeq \frac{m_r^3}{192\pi m_{Pl}^2} \sim \frac{\sqrt{3}M_c^6}{64\pi m_{Pl}^5}, \quad (2.13)$$

where we have ignored a similar contribution from the decay into zero-mode gravitons. For larger M_c , decays into other SM fields such as light fermions and hadrons can increase the width by a factor of order unity. The radion is effectively stable provided τ is larger than $H_0^{-1} \sim 4 \times 10^{17}$ s, corresponding to

$$M_c \lesssim 7 \times 10^8 \text{GeV}. \quad (2.14)$$

2.4 Radion as the Inflaton

The radion potential at leading order is $V_\phi(\phi) \sim m_r^2 \phi^2/2$, for small ϕ/m_{Pl} . At large ϕ/m_{Pl} this form represents an assumption about the form of the $V_\phi(\phi)$. In order for the radion to be the inflaton, the potential energy density of the radion must dominate the energy density of the universe. For the radion to be a viable inflaton candidate, it must satisfy the slow-roll condition. The quadratic potential is of the *chaotic* type, so the slow-roll condition does not depend on the radion mass, but only on the amplitude of ϕ . The slow roll condition can be stated as a condition on a parameter ϵ :

$$\epsilon = \frac{1}{2}m_{Pl}^2 \left(\frac{V'}{V} \right)^2 = 2 \frac{m_{Pl}^2}{\phi^2} < 1, \quad (2.15)$$




Figure 1: Reheat temperature associated with radion decay as a function of the compactification scale.

which is satisfied for $\phi \gtrsim \sqrt{2}m_{Pl}$. As discussed in the next section, this condition turns out to be natural for the radion. Note that the requirement $\phi \gtrsim m_{Pl}$ is still consistent with a regime where our effective theory is under control. Indeed, the mass of the radion being typically much smaller than m_{Pl} , the energy density associated with the radion is still much smaller than m_{Pl}^4 , and therefore we are still away from a regime where 4d quantum gravity corrections would dominate.

One important consideration on inflation is the requirement that the inflaton energy density be converted to radiation after inflation and prior to nucleosynthesis and/or the freeze-out of dark matter relics. The reheating temperature in a scenario with perturbative reheating is given by $T_{RH} \sim \sqrt{\Gamma m_{Pl}} \sim M_c^3/m_{Pl}^2$. We present in Fig. 1 the plot of the decay temperature of the radion as a function of M_c , (*i.e.*, the reheating temperature of the universe in the scenario where the radion is the inflaton). Requiring that the reheating temperature is larger than the TeV scale (so that the electroweak candidates for cold dark matter such as LSPs and/or LKPs are preserved) leads to the requirement $M_c \gtrsim 10^{13}$ GeV.

The fact that the radion couples gravitationally to any field, and in particular to other scalars of the theory, modifies the naïve radion scalar potential written above. The effect is not negligible if the energy of these scalar fields is significant and leads to an interesting inflationary cosmology as we discuss now.

3. Radion Mixing and Induced Inflationary Potential

Let us assume the existence of a scalar field ψ other than the radion. If ψ is a bulk field, according to Eq. (2.7), the total scalar potential is

$$V(\phi, \psi) = V_\phi(\phi) + e^{-\phi/\sqrt{3}m_{Pl}} V_\psi(\psi). \quad (3.1)$$

We continue to assume that the form $V_\phi(\phi) = m_r^2 \phi^2/2$ remains valid at high energies. This provides the simplest phenomenological description of stabilized extra dimensions during inflation. Even if the position of the minimum is displaced because of the second term in Eq. (3.1) and thus the size of the extra dimension is modified, $V_\phi(\phi) \sim m_r^2 \phi^2$ guarantees that the VEV of ϕ remains finite and is not destabilized during inflation. This potential is plotted as a function of ϕ for fixed ψ and a few choices of r in Fig. 2.

When ψ is localized on the 4-d brane, the potential is of a slightly different form:

$$V(\phi, \psi) = V_\phi(\phi) + A e^{-2\phi/\sqrt{3}m_{Pl}} V_\psi(\psi). \quad (3.2)$$

In this case the coupling between the radion and ψ comes from $\sqrt{g}_{\text{induced}} T^\mu{}_\mu$. Here A is an affine function of ξ , where ξ reflects the presence of terms in the (brane) action of the type

$$\sqrt{g}_{\text{induced}} \xi \mathcal{R} (\psi^2 + \dots) \quad (3.3)$$

which mixes the Ricci scalar with ψ^2 . One can suppress the $\phi - \psi$ coupling at tree level by suitably choosing ξ , however this coupling will generically reappear at loop level. Consequently, we expect a potential of the type in Eq. (3.1), and the bulk and brane cases should not be qualitatively very different. We will focus on the bulk case below.

We now investigate the virtues of the potential of Eq. (3.1) for inflation. Because of the $\phi - \psi$ coupling, the VEV of ϕ is shifted from zero and satisfies the relation

$$\langle \phi \rangle = \frac{V_\psi}{\sqrt{3}m_r^2} e^{-\langle \phi \rangle/\sqrt{3}}, \quad (3.4)$$

where from now on, we refer to ϕ (and ψ) in Planck units. We can easily evaluate $\langle \phi \rangle$ in the case where V_ψ is a constant; for instance, in the regime where ψ is slowly rolling. We define the scale M_I to be $V_\psi(\psi) \equiv M_I^4$, and in terms of the parameter r defined as

$$r \equiv \frac{M_c}{M_I}, \quad (3.5)$$

we can express $\langle \phi \rangle$ as

$$\langle \phi \rangle = \frac{e^{-\langle \phi \rangle/\sqrt{3}}}{3\sqrt{3}r^4} \quad \text{and} \quad V(\langle \phi \rangle) = 3M_I^4 r^4 \left(\frac{\langle \phi \rangle^2}{2} + \sqrt{3}\langle \phi \rangle \right). \quad (3.6)$$

We plot in Fig. 3 the VEV of ϕ as a function of r under these conditions.




Figure 2: The scalar potential as a function of ϕ (in Planck units) for $r = 10^1, 10^0, 10^{-1}, 10^{-2}, 10^{-3}, 10^{-4}$ (from left to right). r is defined in Eq. 3.5.

We can see that in the limit $r \gg 1$, the position of the high energy minimum nearly coincides with the position at low energy ($\phi = 0$). When $r \gg 1$ we have $\langle \phi \rangle \sim 1/(3\sqrt{3}r^4)$ leading to $V(\langle \phi \rangle) \approx M_I^4$. Therefore, we recover $\langle \phi \rangle \approx 0$ and the $\phi - \psi$ coupling is irrelevant. As can be seen in Fig. 4, when $r \gg 1$, ψ dominates the potential and plays the role of the inflaton.

In the opposite limit, $r \ll 1$, we find $\langle \phi \rangle \gg 1$ (see Fig. 3), which leads to a suppression of the contribution of ψ in the potential. When $r \ll 1$, ϕ dominates the potential, $V(\langle \phi \rangle) \approx 3M_I^4 r^4 \langle \phi \rangle^2/2$, and ϕ plays the role of the inflaton even though $V_\psi \gg V_\phi$.

A very interesting property of this potential is that even if initially the conditions were not satisfied for ϕ to be an inflaton, ϕ will be driven to be an inflaton through its interactions with ψ . If ψ carries enough energy density, it inevitably pushes the radion away from $\phi = 0$ to large values of ϕ , where the radion can act as an inflaton. Somewhat counter-intuitively, the higher the energy density of ψ , the more efficiently the radion becomes an inflaton. Because of the exponential coupling between ϕ and ψ , the higher M_I , the larger the shift of ϕ , and the larger the eventual suppression of the energy density of ψ . Note that during inflation the size of the extra dimension is modified. It is larger by a factor $e^{\langle \phi \rangle/3}$ compared to the low




Figure 3: The vacuum expectation value of ϕ (in Planck units) at the minimum of its potential as a function of r (as defined in Eq. 3.5). The dashed line corresponds to the case where the inflaton lives in the bulk. The plain lines correspond to cases where the inflaton is localized in four dimensions with suppressed coupling to the radion parametrized by A as defined in Eq. (3.2). From top to bottom, the solid curves are for $A = 0.1, 0.01, 0.001$.

energy case. The mass of the radion is also modified to

$$\frac{\partial^2 V}{\partial \phi^2} = m_r^2 + \frac{V_\psi}{3m_{\text{Pl}}^2} e^{-\phi/\sqrt{3}} \rightarrow m_{\text{eff}}^2 = m_r^2 \left(1 + \frac{\langle \phi \rangle}{\sqrt{3}} \right). \quad (3.7)$$

It is informative to compute the value of the ratio m_{eff}^2/H^2 during inflation:

$$\frac{m_{\text{eff}}^2}{H^2} = \frac{6}{\langle \phi \rangle^2} \frac{1 + \langle \phi \rangle / \sqrt{3}}{1 + 2\sqrt{3}/\langle \phi \rangle}. \quad (3.8)$$

In the limit $r \ll 1$, $m_{\text{eff}}^2/H^2 \rightarrow 2\sqrt{3}/\langle \phi \rangle \ll 1$. As is well known, in this case the primordial fluctuations of ϕ are frozen during inflation. This confirms the fact that the radion is a viable inflaton candidate when $r \ll 1$ since it can be the origin of cosmological perturbations. However, in the opposite limit $r \gg 1$, the effective radion mass is too large: $m_{\text{eff}}^2/H^2 \rightarrow \sqrt{3}/\langle \phi \rangle \gg 1$.




Figure 4: Scalar potential (in units of M_I^4) evaluated at the minimum of ϕ (solid curve). The dashed curve is the contribution from V_ϕ only. The dotted curve is the contribution from ψ : $V_\psi e^{-\langle\phi\rangle/\sqrt{3}m_{Pl}}$. The potential involving the field ψ dominates the energy density for $r \gg 1$, while for $r \ll 1$, ϕ dominates. r is defined in Eq. 3.5.

4. Evolution of the Radion Field

4.1 Evolution During Inflation

We are now interested in determining the evolution of the radion field in the primordial stages of the universe. From the discussion in the previous section, one cannot ignore the coupling between ϕ and ψ . If we want to follow the cosmological evolution of the radion, it is necessary to solve the system of coupled differential equations involving ϕ and ψ . However, for pedagogical purposes, we start with the textbook case and first solve the equation of motion for the VEV of ϕ in the absence of such a coupling :

$$\ddot{\phi} + 3H\dot{\phi} + V'_\phi(\phi) = 0, \quad (4.1)$$

where $H = M_I^2/\sqrt{3}m_{Pl}$ is a constant during inflation. In terms of $\tilde{t} = Ht$ and $x = m_r/H = 3r^2$, the equation of motion reads

$$\ddot{\phi} + 3\dot{\phi} + x^2\phi = 0, \quad (4.2)$$

which has solution

$$\phi(\tilde{t}) = \frac{e^{-3\tilde{t}/2}}{2\Delta} \left[(3\phi_0 + 2\phi_1) \left(e^{\Delta\tilde{t}/2} - e^{-\Delta\tilde{t}/2} \right) + \Delta\phi_0 \left(e^{\Delta\tilde{t}/2} + e^{-\Delta\tilde{t}/2} \right) \right], \quad (4.3)$$

where $\Delta = \sqrt{9 - 4x^2}$, $\phi_0 = \phi(\tilde{t} = 0)$, $\phi_1 = \dot{\phi}(\tilde{t} = 0)$. We typically expect $\phi_1 \sim x\phi_0$ (assuming that the initial energy density of ϕ is equally distributed between kinetic and potential energy). As is well known, for $x \gg 1$, ϕ is exponentially damped during inflation while for $x \ll 1$, ϕ is frozen and remains unaffected by inflation. However, as we will see now, these conclusions are substantially altered when including a coupling between the moduli and the inflaton, even as weak as a gravitational coupling.

If ψ is a bulk field, the coupled equations to be solved are

$$\begin{aligned} \ddot{\phi} + \sqrt{3}\dot{\phi} \left\{ \frac{\dot{\phi}^2}{2} + \frac{\dot{\psi}^2}{2} + c_r \frac{\phi^2}{2} + \frac{V_\psi e^{-\phi/\sqrt{3}}}{\mathcal{M}^2 m_{Pl}^2} \right\}^{1/2} + d_r \dot{\phi} + c_r \phi - \frac{V_\psi e^{-\phi/\sqrt{3}}}{\sqrt{3}\mathcal{M}^2 m_{Pl}^2} &= 0 \\ \ddot{\psi} + \sqrt{3}\dot{\psi} \left\{ \frac{\dot{\phi}^2}{2} + \frac{\dot{\psi}^2}{2} + c_r \frac{\phi^2}{2} + \frac{V_\psi e^{-\phi/\sqrt{3}}}{\mathcal{M}^2 m_{Pl}^2} \right\}^{1/2} + d_i \dot{\psi} + \frac{V'_\psi(\psi) e^{-\phi/\sqrt{3}}}{m_{Pl} \mathcal{M}^2} &= 0, \end{aligned} \quad (4.4)$$

where ϕ and ψ are in Planck units, we have defined a dimensionless time variable $\tilde{t}$ in terms of a convenient mass scale $\mathcal{M}$ as $\tilde{t} = \mathcal{M}t$, and we have introduced the constants

$$c_r = \frac{m_r^2}{\mathcal{M}^2}, \quad d_r = \frac{\Gamma_\phi}{\mathcal{M}}, \quad d_i = \frac{\Gamma_\psi}{\mathcal{M}}. \quad (4.5)$$

Note that the addition of the decay rate terms in the equations of motion is a an effective description which is valid only when oscillations of the fields have started [24].

We define D to be the value of ϕ in Planck units when the field oscillates and its energy density starts to decrease as a nonrelativistic component:

$$D \equiv \frac{\phi(t_{osc})}{m_{Pl}}. \quad (4.6)$$

The factor D may be interpreted as the damping factor if the initial value of the radion is Planckian. Our goal is to evaluate by which amount the radion field can be damped during inflation for a given value of r , so that we can investigate the resulting radion energy density today.

The damping of the amplitude of ϕ generally depends on the form of the potential for ψ . However, some general statements can be made. Indeed, what typically happens can be easily understood by inspection of Fig. 3: Because of the coupling between ϕ and ψ , the position of the minimum for ϕ is displaced from its low energy position. During inflation, ϕ is pushed to its high-energy minimum and we naturally expect that once inflation ends, oscillations in ϕ start with an amplitude of the order of the VEV of ϕ as given in Fig. 3. We checked that this is indeed what typically happens using different types of potentials for ψ .




Figure 5: Examples of the time evolution of ϕ and ψ using the potential for ψ of the form $V_\psi(\psi) = m_\psi^2 \psi^2/2$. In these specific examples, $r = 0.5$ in the top panel and $r = 10$ in the bottom panel ($r = \sqrt{m_r/m_\psi}$). Initial conditions were chosen such that $\psi_0 = 5$, and ϕ_0 to satisfy the equipartition condition.

As an illustration, we show in Fig. 5 the time evolution for the ϕ and ψ fields in the case where the potential for ψ is of the large-field (chaotic inflation) type.² What is shown in the

²An interesting possibility for the origin of the “precursor” inflaton ψ is A_5 , the higher component of a gauge field [25].




Figure 6: $D(r)$ as derived in the chaotic case. For initial conditions we have assumed $\psi_0 = 5$, and have chosen ϕ_0 to satisfy the equipartition condition.

figures are the values of $\langle\phi^2\rangle^{1/2}$ and $\langle\psi^2\rangle^{1/2}$:

$$\langle\phi^2\rangle^{1/2} \equiv \sqrt{\rho_\phi/m_r^2} \quad (4.7)$$

$$\langle\psi^2\rangle^{1/2} \equiv \sqrt{\rho_\psi/m_\psi^2 \exp(-\phi/\sqrt{3})}, \quad (4.8)$$

where the energy densities capture both the kinetic and potential energies. During inflation, $\langle\phi^2\rangle^{1/2} = \phi/\sqrt{2}$ and $\langle\psi^2\rangle^{1/2} = \psi e^{-\phi/2\sqrt{3}}/2$. After inflation, when the fields oscillate, the averages defined above correspond to cycle averages of the field.

Note that in the case of the chaotic potential $m_\psi^2\psi^2/2$, $r \sim \sqrt{m_r/m_\psi}$. For r to be larger than unity would require that the mass of ψ be somehow less than the radion itself. This may be difficult to achieve naturally, however, we consider this possibility since we find that the chaotic potential for ψ provides the largest damping during inflation. (See the $D(r)$ plot derived in this case in Fig. 6). As initial conditions, one may choose equipartition of the potential energy densities between ϕ and ψ , leading to

$$e^{-\phi_0/\sqrt{3}} = r^4 \phi_0^2/\psi_0^2, \quad (4.9)$$

and the initial condition for ψ_0 has to be chosen such that ψ_0 satisfies a slow-roll condition.

In the following sections we will express cosmological constraints as general bounds on D , and will illustrate how these constraints typically translate into r , remembering that for $r < 1$, we expect D will naturally be of order unity.

4.2 Validity of 4D effective field theory

Throughout the discussion we have assumed an effective 4-dimensional description. The 4d effective field theory is an approximation which neglects the effects of higher KK modes. Those can be excited at high temperature. For the 4d description to be valid, we should have the temperature smaller than M_c . This requirement is in addition to the 5d cut-off Λ which characterizes the scale at which the 5d gauge theory (including KK modes) becomes strongly coupled, and loses predictivity.

The definition of the frontier between the 4d and 5d descriptions in the inflationary phase, where the universe is dominated by vacuum energy, is somewhat subtle. During inflation it is possible to define a temperature. In the exponentially expanding universe particles are created with typical momentum of the order of H . In this phase, the criterion for the validity of the 4d effective description is $H \lesssim L_{phys}^{-1}$, where $L_{phys} = e^{\langle\phi\rangle/\sqrt{3}}/M_c$. This leads to a constraint on M_c , shown in Fig. 7. The corresponding constraint is weak, and is only relevant for small values of r , which, as we will see in the next section, will be disfavored anyway because of the problem of the overclosure of the universe by radions.

The reheat temperature satisfies $T_{RH} \lesssim M_I$ where the bound is saturated in the case of preheating. We can define a parameter $\mathcal{V} \equiv T_{RH}/M_I$ which measures the efficiency of the transfer between the inflaton energy to radiation energy. Requiring an effective 4d description ($T_{RH} \lesssim M_c$) leads to $\mathcal{V} \lesssim r$. Note that in the case of perturbative reheating $T_{RH} \sim \lambda\sqrt{m_I m_{Pl}}$ where λ is the inflaton coupling to normal matter and m_I is the inflaton mass. The requirement that the theory appear four dimensional thus translates into $m_I \lesssim r^2 \lambda^{-2} 10^{11}$ GeV.

5. Radion as Dark Matter

5.1 Coherent Production

We now turn to the evolution of the radion field once inflation is completed. The story begins with the energy density of the radion after inflation (determined by the parameter D discussed in the previous section). From the time oscillations commence until today,³ the radion energy density ρ_r redshifts as nonrelativistic matter, so that

$$\rho_r(t_0) = \rho_r(t_{osc}) \frac{a_{osc}^3}{a_0^3}, \quad (5.1)$$

where the subscript ‘0’ refers to the present, and

$$\rho_r(t_{osc}) = \frac{1}{2} m_r^2 \phi^2(t_{osc}) = \frac{3}{2} D^2 M_c^4. \quad (5.2)$$

³If the radion is dark matter it must have a lifetime longer than the age of the universe, which requires $M_c \lesssim 7 \times 10^8$ GeV.




Figure 7: For the Hubble radius to be larger than the size of the extra dimension $L_{phys} = e^{\phi/\sqrt{3}}/M_c$ during inflation, M_c must be in the lower region limited by the dashed line. This does not lead to strong constraints in the preferred region $r \gtrsim 1$.

After inflation, until reheating (or preheating), the energy densities in both the inflaton and the radion decrease as a nonrelativistic component. We can express $\rho_r(t_0)$ as

$$\rho_r(t_0) = \frac{3}{2} D^2 M_c^4 \frac{a_{osc}^3}{a_{RH}^3} \frac{a_{RH}^3}{a_0^3}. \quad (5.3)$$

Before reheating, $a^3 \propto t^2$. Since $t^2 \sim H^{-2} \propto \rho^{-1}$, we can replace the factor a_{osc}^3/a_{RH}^3 by ρ_{RH}/ρ_{osc} . The value of ρ_{RH} is simply $(\pi^2 g_*(T_{RH})/30) T_{RH}^4$. The value of ρ_{osc} is not so straightforward.

Here we will use the results for the chaotic inflation example of the previous section. If $r \lesssim 1$, then the radion is the inflaton. In this case inflation ends when $\phi \simeq m_{Pl}$ and $\rho_{osc} \simeq m_r^2 m_{Pl}^2$. Also in this case, $D \simeq 10^{-1}$. If $r \gtrsim 1$ the energy during inflation is approximately M_I^4 and $D \simeq 10^{-2} r^{-4}$ (as can be seen from Fig. 6). Using these values, along with the fact that $a_{RH}^3/a_0^3 = g_*(T_0) T_0^3 / g_*(T_{RH}) T_{RH}^3$, we find

$$\rho_r(t_0) = \frac{\pi^2}{20} g_*(T_0) T_0^3 T_{RH} \begin{cases} 3 \times 10^{-3} & r \lesssim 1 \\ 10^{-4} r^{-4} & r \gtrsim 1. \end{cases} \quad (5.4)$$

Using $T_0 = 2.4 \times 10^{-13}$ GeV, $g(T_0) = 3.36$, and $\rho_c = 3.9 \times 10^{-47}$ GeV⁴, this leads to a value of Ω_r of

$$\Omega_r = \begin{cases} 2 \times 10^6 (T_{RH}/\text{GeV}) & r \lesssim 1 \\ 6 \times 10^4 (T_{RH}/\text{GeV}) r^{-4} & r \gtrsim 1. \end{cases} \quad (5.5)$$

If the radion is stable, we require $\Omega_r \lesssim 0.3$, and the inequality is saturated if the radion is dark matter. The limit to Ω_r results in the limit $T_{RH} \leq 5 \times 10^{-7}$ GeV for $r \lesssim 1$, and $T_{RH} \leq 2 \times 10^{-5} r^4$ GeV for $r \gtrsim 1$. Clearly the $r \lesssim 1$ case is ruled out. The parameter r must be sufficiently large to avoid overclosure:

$$r \gtrsim 15 \left(\frac{T_{RH}}{\text{GeV}} \right)^{1/4}. \quad (5.6)$$

We now discuss some potential observational effects associated with the oscillation of the radion⁴. Coherent oscillations of the radion induce variations of gauge couplings (in a way described in Ref. [14], for instance). For this effect not be observable, we require the oscillation period m_r^{-1} to be much smaller than the Hubble time. There are constraints on the variation of the gauge couplings at the time of nucleosynthesis. Applying the above requirement to the nucleosynthesis epoch leads to $H_{nucleo} \ll m_r$, *i.e.*, $T_{nucleo} \ll M_c$, which is always satisfied. Note that as soon as structures start to form, coherent oscillations will get distracted by the development of gravitational potentials. Therefore, radion oscillations would no longer be coherent today and would not be seen as variations of the couplings, but instead, as nonrelativistic matter.

5.2 Thermal Production

Now we turn to the thermal production of radions at reheating. The number density of radions produced thermally is obtained by solving the Boltzmann equation

$$\frac{dn_r}{dt} + 3Hn_r = -\langle\sigma v\rangle n_\gamma (n_r - n_r^{eq}) - \Gamma(n_r - n_r^{eq}), \quad (5.7)$$

where n_γ represents the number density of a typical light species in the reaction $\text{radion} + \gamma \rightarrow \text{everything}$. The cross section is expected to be approximately m_{Pl}^{-2} , and the decay width is of order m_r^3/m_{Pl}^2 .⁵ The feeble radion interactions with matter imply that the radion has great difficulty reaching thermal equilibrium. We have also seen above that the coherent production of radions need be very small. Thus, initially, $n \ll n_{eq}$. In terms of $Y \equiv n_r/s$ where s is the entropy density, and $x \equiv m_r/T$, the Boltzmann equation becomes

$$\frac{dY}{dx} = \frac{m_r}{m_{Pl}} \left(\frac{C_A}{x^2} + x C_D \right) Y_{eq}, \quad (5.8)$$

where C_A and C_D are dimensionless numbers.

⁴See also [16, 26]

⁵There are potentially large dimensionless numbers in the cross section and decay width; *e.g.*, the factor of 192π appearing in the decay width given in Eq. (2.13).

If the radions are relativistic ($x \ll 1$), $Y_{eq} \sim x^{-3}$, and

$$Y \sim \frac{T_{RH}}{m_{Pl}}. \quad (5.9)$$

We note again that there is a potentially small (say 10^{-2}) dimensionless number on the right-hand-side of the above equation.

We now examine the case where the radions are nonrelativistic at the time of reheating ($x \gg 1$) so that $Y_{eq} \sim x^{3/2} \exp(-x)$ and we can ignore the first term in the right hand side of Eq. (5.8). In this case

$$Y \sim \frac{m_r}{m_{Pl}} \mathcal{I}(x_{RH}), \quad (5.10)$$

with

$$\mathcal{I}(x_{RH}) \equiv \int_{x_{RH}}^{\infty} x^{5/2} e^{-x} dx, \quad (5.11)$$

where $x_{RH} = m_r/T_{RH}$ is x at the time of reheating.

We find that Ω_r from thermal radions (for relativistic radion production) is

$$\Omega_r = 10^9 \frac{T_{RH}}{\text{GeV}} \left(\frac{M_c}{m_{Pl}} \right)^2, \quad (5.12)$$

while the value for nonrelativistic radion production is

$$\Omega_r = 10^9 \text{GeV}^{-1} \frac{M_c^4}{m_{Pl}^3} \times \mathcal{I}(x_{RH}). \quad (5.13)$$

The limit $\Omega_r < 0.3$ becomes

$$\left(\frac{M_c}{5 \times 10^{13} \text{GeV}} \right)^2 \frac{T_{RH}}{\text{GeV}} < 1 \quad (5.14)$$

for relativistic radion production, and for nonrelativistic radion production

$$\left(\frac{M_c}{4 \times 10^{11} \text{GeV}} \right)^4 \mathcal{I}(x_{RH}) < 1. \quad (5.15)$$

This last bound is always satisfied, while the bound from relativistic radions only leads to a very weak constraint on the reheat temperature. Therefore, there is no consequence, as far as dark matter is concerned, from thermally produced radions.

Radionactive dark matter would be challenging to detect directly, though it may have implications either for gamma-ray spectra (through the eventual decay of the radions) or isocurvature perturbations.

6. Radions and Cosmological Perturbations

6.1 Radion Isocurvature Perturbations

A possible effect from the stable radion is on the CMB, an issue we now investigate. Indeed, in a way reminiscent of the axion, there are isocurvature perturbations associated with the radion as dark matter. To discuss this, we begin with a few definitions. The curvature perturbation associated with an energy density component ρ_i is defined by

$$\zeta_i = -H \frac{\delta \rho_i}{\dot{\rho}_i}. \quad (6.1)$$

For matter, $\zeta_m = \frac{1}{3} \delta \rho_m / \rho_m$ and for radiation $\zeta_\gamma = \frac{1}{4} \delta \rho_\gamma / \rho_\gamma$. The adiabatic mode of the perturbation is defined by $\zeta_i = \zeta_\gamma$. The nonadiabatic or isocurvature modes arise from local inhomogeneities in the equation of state of the energy density. They may be specified relative to ζ_γ as

$$\mathcal{S}_i \equiv \frac{\delta \rho_i}{\rho_i} - \frac{3}{4} \frac{\delta \rho_\gamma}{\rho_\gamma} = 3(\zeta_i - \zeta_\gamma) \quad (6.2)$$

The large-scale temperature anisotropy can be related to the primordial adiabatic and isocurvature perturbations by (see for instance section II of Ref. [27]):

$$\frac{\delta T}{T} \approx -\frac{1}{5} \zeta^{ad} - \frac{2}{5} \left(\frac{\rho_c}{\rho_m} \mathcal{S}_c + \frac{\rho_b}{\rho_m} \mathcal{S}_b \right) + \frac{1}{15} \frac{\rho_\nu}{\rho_\gamma + \rho_\nu} \mathcal{S}_\nu \quad (6.3)$$

where ζ^{ad} measures the primordial adiabatic perturbation, $\mathcal{S}_c$, $\mathcal{S}_b$ and $\mathcal{S}_\nu$ the isocurvature perturbations in cold dark matter, baryons and neutrinos respectively. Here ρ_m is the total matter energy density, $\rho_m = \rho_c + \rho_b$.

We are interested in constraining $\mathcal{S}_c$. From an observational point of view, what is constrained is the quantity:

$$B = \frac{\mathcal{S}_b^{\text{eff}}}{\zeta^{ad}}, \quad (6.4)$$

where

$$\mathcal{S}_b^{\text{eff}} = \mathcal{S}_b + \frac{\rho_c}{\rho_b} \mathcal{S}_c \quad (6.5)$$

is the effective nonrelativistic term. We use the constraints derived at 95% C.L. in Ref. [27]

$$-0.46 < B < 0.35. \quad (6.6)$$

In the case where the radion is the dark matter we have

$$\mathcal{S}_c = \mathcal{S}_{\text{radion}} = 3(\zeta_{\text{radion}} - \zeta_\gamma) = \frac{\delta \rho_\phi}{\rho_\phi} = \delta^{iso}, \quad (6.7)$$

where, at first order,

$$\delta^{iso} = 2 \frac{\delta \phi}{\phi}. \quad (6.8)$$

If the radion is practically free during inflation, with small effective mass $m^2 \ll H^2$, it follows that on superhorizon scales

$$\frac{\delta\phi}{\phi} \approx \frac{H_*}{2\pi\phi_*}, \quad (6.9)$$

where H_* and ϕ_* are H and ϕ evaluated at the time of horizon exit, and

$$\delta^{iso} = \frac{H_*}{\pi\phi_*} = \frac{\sqrt{V_*}}{\sqrt{3}\pi D_* m_{Pl}^2}, \quad (6.10)$$

where D_* is ϕ_*/m_{Pl} . Equation (6.6) leads to the restriction $|\delta^{iso}| \lesssim 10^{-2}\zeta_\gamma$, which leads to

$$\sqrt{V_*} \lesssim 10^{-5} D_* m_{Pl}^2. \quad (6.11)$$

This constraint, while interesting in its own right, is not relevant for our current picture. As we will see in the next section, $m^2/H^2 \lesssim 1$ during inflation only if $r \lesssim 0.4$, which typically results in D of order unity. But if D is of order unity, the limit in Eq. (6.11) is weaker than the limit on V_* from the limit to the gravitational wave background produced in inflation. However, we note that there may exist unconventional inflationary mechanisms where this limit is useful.

To summarize, we have seen that when the compactification scale is below 7×10^8 GeV and the radion is effectively stable, to avoid overclosure of the universe by radions we must have $r \gtrsim 15(T_{RH}/\text{GeV})^{1/4}$.

If D and r are properly correlated, radions could in fact play the role of cold dark matter, though in the models of inflation we have considered this would require fine tuning.

6.2 Radion as the Curvaton

It is now widely believed that the origin of structure in the universe is a primordial perturbation already in existence when cosmological scales start to enter the horizon. In the standard picture for the generation of cosmological perturbations, the curvature perturbation ζ is generated during inflation through the perturbation of a single component inflaton.

In the curvaton scenario [11] the curvature perturbation produced during inflation is initially negligible, but the curvaton field acquires “isocurvature” perturbations during inflation, which is converted to curvature perturbations when the curvaton density becomes a significant fraction of the total density. In this case, ζ is given by

$$\zeta = f_D \times \frac{\delta^{iso}}{3} \quad (6.12)$$

where f_D is the fraction of the total energy density (radion r and radiation R) of the universe in the radion–curvaton when it decays:

$$f_D = \left[\frac{\rho_r}{\rho_r + \frac{4}{3}\rho_R} \right]_{H=\Gamma}. \quad (6.13)$$

When the radion decays, it reheats the universe and imprints its perturbation into photons and matter. The degree of nonadiabaticity is given by the fraction f_D which reflects how much of the total energy density is carried by the radion at the time of its decay. If this fraction f_D is unity, there will be no residual isocurvature perturbations. On the other hand, if the radion does not completely dominate the energy density of the universe at the time of decay, there will be residual isocurvature perturbations.

If $m^2 \ll H^2$ during inflation then the spectrum of the fractional field perturbation is

$$\mathcal{P}_{\delta\phi/\phi}^{1/2} \approx \frac{H_*}{2\pi\phi_*} \quad (6.14)$$

and that of the fractional energy density perturbation is

$$\mathcal{P}_{\delta^{iso}}^{1/2} \approx \frac{H_*}{\pi\phi_*} \approx \delta^{iso} \quad . \quad (6.15)$$

It follows that the prediction of the radion-curvaton model for the spectrum of the curvature perturbation is

$$\mathcal{P}_\zeta^{1/2} \approx \frac{f_D}{3} \delta^{iso} \approx \frac{f_D H_*}{3\pi\phi_*}. \quad (6.16)$$

This is to be compared with the COBE measurement of the CMB quadrupole anisotropy

$$\mathcal{P}_\zeta^{1/2} = 4.8 \times 10^{-5} \quad . \quad (6.17)$$

It is now clear that the radion cannot easily play the role of a curvaton. Indeed, from Fig. 8, we see that the effective mass of the radion is smaller than H only for $r \lesssim 0.4$. On the other hand, Fig. 4 indicates that for small r , ϕ dominates the total energy density during inflation, in conflict with the curvaton scenario. In fact, this leads back to the model of the radion as the inflaton (and providing the primordial fluctuations) discussed in Sec. 2.4.

It is possible that ψ could play the role of the curvaton. This would require that ψ decays after ϕ , which may be unlikely for low compactifications scales since generically a scalar field would have gravitational interactions, and thus at most a lifetime of the same order as the radion. This would require $\lambda m_\psi \lesssim M_c^6/m_{Pl}^5$, where λ characterizes the ψ interaction strength.

Let us now see if the radion can play the role of a *modulon*.

6.3 Radion as the Modulon

As mentioned in the introduction, another alternative to the standard picture of generation of cosmological fluctuations was recently proposed [12], which also makes use of a light scalar field that controls the coupling of the inflaton to matter. To distinguish it from the curvaton scenario, we refer to the field responsible for the generation of fluctuations in this mechanism as a *modulon*. In the modulon scenario, the coupling λ between the inflaton and ordinary matter is not constant but is given by the fluctuations around the VEV of the modulon. If the modulon is light enough, it acquires primordial fluctuations during inflation of order $H_*/2\pi$. These spatial fluctuations will then correspond to spatial fluctuations in the coupling λ . In the




Figure 8: The ratio m_{eff}^2/H^2 during inflation (evaluated at the high-energy minimum of the potential for ϕ). Fluctuations in ϕ are frozen and can be at the origin of cosmological perturbations only for $r \lesssim 0.4$.

following we will denote the modulon by ϕ since we want to check whether the radion can be a modulon candidate. The (local) reheat temperature depends on this coupling according to

$$T_{RH} \sim \sqrt{\Gamma_I m_{Pl}}, \quad (6.18)$$

where $\Gamma_I \sim \lambda^2 m_I$ is the decay rate of the inflaton and m_I its mass. In the modulon scenario,

$$\frac{\delta T}{T} \sim \frac{\delta \lambda}{\lambda}, \quad (6.19)$$

where λ is given by

$$\lambda(\phi) = \lambda_0 \left(1 + \frac{\phi}{M} + \dots\right). \quad (6.20)$$

As the radion controls the size of the extra dimension, it controls the inflaton coupling to matter as in Eq. (6.20), where λ_0 is related to a dimensionful, 5d parameter by $\lambda_0 = \lambda_5/L$. Small variations in the radion VEV thus indeed lead to variations in the coupling between matter and the inflaton, with $M = \sqrt{3}m_{Pl}$. The fraction of the coupling controlled by the fluctuating radion is ϕ_*/m_{Pl} where $\phi_* = D_* m_{Pl}$ is the value of ϕ at the time of horizon exit so that Eq. (6.19) can be written as

$$\frac{\delta T}{T} \sim \frac{\phi_*}{m_{Pl}} \frac{H_*}{2\pi\phi_*} \sim \frac{H_*}{2\pi m_{Pl}}, \quad (6.21)$$

a contribution typically much too small to account for observed inhomogeneities. Thus, in models with flat extra dimensions, the radion cannot effectively play the role of a modulon. For this mechanism to be effective, the radion coupling should be much larger, as for example in warped compactifications. In the Randall-Sundrum geometry, we would arrive at $\delta T/T \sim H_*/1$ TeV. In order to explain the COBE result, this would require an inflation scale around 10^8 GeV, which is neither the fundamental UV nor the IR scale of the Randall-Sundrum model, but could be generated by an intermediate brane [28].

7. Late Decay of Radions and KK Gravitons

The decay of massive relics after Big Bang Nucleosynthesis produces energetic photons which may dissociate light elements and modify the predictions of their abundances. The constraint on the decay of heavy relics from BBN is (see for instance [29])

$$mY \lesssim 10^{-12} \text{ GeV} \quad (7.1)$$

where $Y \equiv n_X/s$ is the ratio we would have observed today if particle *had not decayed*, and $s \sim T^3$ is the entropy per comoving volume. This bound literally applies for particles whose lifetime is 10^8 s; bounds for larger lifetimes are somewhat relaxed. Thus, by taking 10^{-12} GeV on the right hand side of Eq. (7.1) we are making a conservative estimate for the allowed parameter space. Note that the constraint Eq. (7.1) applies only if the decay takes place after nucleosynthesis.

7.1 Late Decay of Radions

From our estimate of the radion lifetime, Eq. (2.13), we find that the radion decays after nucleosynthesis ($\tau \gtrsim 10^{-2}$; $\Gamma \lesssim 10^{-22}$ GeV) provided

$$7 \times 10^8 \text{ GeV} \lesssim M_c \lesssim 10^{12} \text{ GeV}, \quad (7.2)$$

where the lower limit on M_c comes from the requirement that the radion lifetime is smaller than the age of the universe. In order to explore the implications on radion cosmology, we now estimate $Y_r = n_r/s$ at the time of reheating (which is the same as the ratio we would have observed today if the expansion is adiabatic).

The bound in Eq. (7.1) may be written as

$$m_r \frac{n_r}{s} = \frac{\rho_r}{s} = \Omega_r \frac{\rho_c}{s} \leq 10^{-12} \text{ GeV}. \quad (7.3)$$

Using $\rho_c/s \sim 10^{-9}$, we obtain the constraint $\Omega_r \lesssim 10^{-4}$.

For the chaotic inflationary potential, coherent radion production [using Eq. (5.5)] yields

$$\frac{T_{RH}}{\text{GeV}} \lesssim \begin{cases} 5 \times 10^{-11} & r \lesssim 1 \\ 2 \times 10^{-9} r^4 & r \gtrsim 1. \end{cases} \quad (7.4)$$

Recall that this bound only applies in the range $7 \times 10^8 \text{ GeV} \lesssim M_c \lesssim 10^{12} \text{ GeV}$, so that the radion decays after nucleosynthesis. Obviously the case $r \lesssim 1$ is again excluded. The bound on T_{RH} for $r \gtrsim 1$ is plotted on figure 9.

The constraint in Eq. (7.1) also applies to radions of thermal origin:

$$T_{RH} \lesssim \left(\frac{10^{12} \text{ GeV}}{M_c} \right)^2 \text{ GeV} \quad (7.5)$$

from the thermal production of relativistic radions in the radiation era. If $M_c \sim 10^9 \text{ GeV}$, then $T_{RH} \lesssim 10^6 \text{ GeV}$. If $M_c \sim 10^{12} \text{ GeV}$, then the bound becomes $T_{RH} \lesssim 1 \text{ GeV}$.

For nonrelativistic radion production [using Eq. (5.13)] the bound in Eq. (7.1) becomes

$$\mathcal{I}(x_{RH}) \lesssim \left(\frac{4 \times 10^{10} \text{ GeV}}{M_c} \right)^4. \quad (7.6)$$

This bound imposes constraints on T_{RH} for the larger region of M_c . Below $M_c \lesssim 3 \times 10^{10} \text{ GeV}$, there is no bound on T_{RH} . At $M_c \sim 10^{12} \text{ GeV}$, the bound is $T_{RH} \lesssim 35 \text{ TeV}$, for $M_c \sim 10^{11} \text{ GeV}$, the bound is $T_{RH} \lesssim 800 \text{ GeV}$ and for $M_c \sim 5 \times 10^{10} \text{ GeV}$, it is $T_{RH} \lesssim 320 \text{ GeV}$.

7.2 Late Decay of KK gravitons

KK gravitons (G^1) also couple very weakly, $\sim m_{Pl}^{-1}$. They have a mass characterized by the size of the extra dimension, $m_{KK} \sim M_c$ and therefore a decay width of the order $\Gamma \sim M_c^3/m_{Pl}^2$, though the specific details of the expression depend on whether KK parity is well conserved, which would forbid decays directly into zero modes of SM fields. Continuing with dimensional analysis, the decay is after nucleosynthesis if $M_c \lesssim 8 \times 10^4 \text{ GeV}$. On the other hand, we only consider values of M_c above several hundred GeV, as lower values are generally excluded by collider constraints [17]. Thus, constraints on KK gravitons from BBN will only apply in the narrow range

$$300 \text{ GeV} \lesssim M_c \lesssim 8 \times 10^4 \text{ GeV} \quad (7.7)$$

In this range of values for M_c , the radion is stable, and to avoid overclosure of the universe by radions, r (D) and T_{RH} must be chosen appropriately in Eq. (5.5).

We assume the universe is radiation dominated at reheating, and compute bounds from the thermal production of KK gravitons. As for the radion, $\langle \sigma v \rangle \sim m_{Pl}^{-2}$, and many of the radion results can be carried over with $m_r \rightarrow M_c$. We begin with the case of relativistic KK gravitons, $T_{RH} \gtrsim M_c$. In this case, thermal production of KK gravitons leads to $Y \sim T_{RH}/m_{Pl}$ and the BBN bound results in,

$$T_{RH} \lesssim \left(\frac{10^6 \text{ GeV}}{M_c} \right) \text{ GeV} \quad (7.8)$$

from the late decay of thermally produced, relativistic KK gravitons. For M_c at the TeV scale, this roughly constrains T_{RH} to be of order M_c . For somewhat smaller extra dimensions, this




Figure 9: The constraints from overclosure by radions (produced non-thermally) –dashed curve– as well as from BBN –solid curve– due to the late decay of radions, translated into a bound on T_{RH} as a function of r . The bound from overclosure (see Eq. 5.5) applies in the range $M_c \sim 10^3 - 7 \times 10^8$ GeV and the one from BBN (Eq. 7.4) in the range $M_c \sim 7 \times 10^8 - 10^{12}$ GeV.

requires a smaller reheating temperature, and indicates that KK gravitons cannot in fact be relativistic for $M_c \gtrsim 1$ TeV.

For nonrelativistic KK gravitons, $Y \sim M_c/m_{Pl}x_{RH}\mathcal{I}$, where now $x_{RH} = M_c/T_{RH}$. Thus, the bound on the reheating temperature is,

$$\mathcal{I} \frac{M_c}{T_{RH}} \lesssim \left(\frac{10^3 \text{ GeV}}{M_c} \right)^2 \quad (7.9)$$

which for all M_c of interest results in a bound of $T_{RH} \lesssim M_c/\text{a few}$ ⁶, consistent with the possibility of KK dark matter [5], and in fact with the general WIMP hypothesis, for which the freeze-out temperature is typically a few tens of GeV. Note also that Ref. [31] studied the case where KK dark matter was made of KK gravitons rather than KK hypercharge gauge bosons, with the relic abundance determined initially by freeze-out of B^1 . They derived bounds from BBN on this scenario coming from the eventual late decay of B^1 into G^1 and a photon.

⁶This point was realized some time ago by Abel and Sarkar [30]

To finish this section, we discuss the possibility of nonthermal production of KK gravitons. As stressed in [32], the gravitational production of particles during inflation can be very efficient and can lead to much stronger bounds on the reheat temperature than the ones from thermal production. However, for this mechanism to be operational, we need the mass of the particle to be smaller than H during inflation. In our case, this corresponds to $e^{-\phi/3}M_c < H$.

From Fig 7, this leads to a bound on M_c only for very small values of r . On the other hand, because of the bound from overclosure of the universe we typically have $r \gtrsim 1$ so that no KK gravitons are expected to be produced nonthermally. Note also that from Fig. 8, there is essentially no gravitational production of radions during inflation for $r \gtrsim 1$. In any case, the bound from overclosure of the universe by radions is not the only reason why a small hierarchy between M_c and M_I is required. A large enough reheat temperature, to have WIMP dark matter, also requires large values of r .

7.3 Diffuse Gamma-ray Signal

Finally, for a very long lived radion, there might be some interesting observable diffuse gamma-ray signal [29, 33]. This applies for life times larger than 10^{14} s, *i.e.*, for

$$M_c \lesssim 3 \times 10^9 \text{ GeV}. \quad (7.10)$$

On the other hand, the radion becomes stable for $\tau \gtrsim 10^{17}$ s, *i.e.*, for $M_c \lesssim 9 \times 10^8 \text{ GeV}$. The observable signal therefore corresponds to a tiny window of values for M_c . This is because the decay rate depends on the sixth power of M_c . Note that there is no similar expected signal from decaying KK gravitons (however see [31] for the special case of UED) since $\tau \gtrsim 10^{17}$ s corresponds to the disallowed region $M_c \lesssim 0.4 \text{ GeV}$.

8. Conclusions

In this work we have studied the cosmological properties of the radion field as it appears in models with flat or almost flat extra dimensions. We find that the potential roles the radion may play in cosmology are very sensitive to the scale (and model) of inflation. Thus, our work explores the challenges involved in building a realistic cosmological model for theories with flat extra dimensions. Figure 10 summarizes our results.

Its origin as a component of the higher dimensional graviton tensor dictates the radion's couplings. Similarly, the mass of the radion, in the absence of fine tuning, is generally related to the compactification scale M_c by $m_r \sim M_c^2/m_{Pl}$. Thus, once one has chosen the size of the extra dimension, the natural radion properties are determined. Of course, it is possible that the radion potential is itself fine tuned by some mechanism we currently do not understand. In that case, its couplings to other fields may still be of gravitational strength, but its mass could possibly differ from our estimates. One avenue of future research would be to explore the compactification dynamics that could naturally make the radion, for example, more massive than the naïve estimates, such that it decays at early times and evades some cosmological constraints.




Figure 10: A summary of the cosmological significance in various regions of radion model parameter space. M_c is the compactification scale and M_I is the scale of the potential of the scalar field responsible for inflation. We note that the application of effective field theory breaks down in the region $M_I \gtrsim 10^3 M_C$, as well as the region where the Hubble radius is smaller than the size of the extra dimension.

We have written the scale of the radion potential as M_c and used a naturalness argument to identify $L \sim M_c^{-1}$, without specifying the compactification dynamics. It is interesting to compare this with existing stabilization models, and the radion masses which result. In Ref. [34], it was shown that casimir effects will stabilize the extra dimension (and in fact, are always present for a compact space). This leads to a potential which is natural in our sense, and an estimate of the radion mass similar to Eq. (2.11) [14]. An alternative (in 6 dimensions) is to invoke a bulk cosmological constant and a set of gravitating 3-branes, balanced against quantum effects from KK mode gravitons [35]. This also leads naturally to the relation between the scale of the potential (characterized, *i.e.*, by the bulk cosmological constant) and L which we have assumed. Unlike the TeV^{-1} (or smaller) extra dimensions we consider, the aim of those works was to generate large (mm) size extra dimensions, for which the radion is too light, in conflict with precision measurements of gravity. In order to generate acceptable large extra-dimension phenomenology, one may introduce a bulk $U(1)$ gauge theory and stabilize with its trapped magnetic flux [35, 36]. An additional parameter allows one to tune the radion potential such that the simple relation between the radion

mass and M_c is disturbed, with the free parameter given by the $U(1)$ coupling. Another way to avoid the natural relation $m_r \sim M_c^2/m_{Pl}$ was suggested in [37] where the stabilizing potential depends only logarithmically on the radion field so that a large hierarchy between M_c and R^{-1} can be generated without fine-tuning. The idea works for an even number of dimensions only since it relies on sets of two transverse dimensions (for the logarithmic dependence). In addition, bulk supersymmetry is needed to protect against the generation of a bulk cosmological constant which would induce power law corrections to the effective potential for the radion and spoil the picture with logarithmic potentials.

Concerning the radion coupling to an inflaton (or any scalar field), the result is an interesting inflationary dynamics through which the radion may be pushed to dominate the energy density of the universe and play the role of the inflaton. Given the scale M_I which characterizes the potential of another scalar field, the cosmological evolution of the universe depends crucially on the ratio $r = M_c/M_I$ where M_c is the compactification scale. The coupling to the inflaton will generally push the radion VEV away from its low energy minimum, in way strongly reminiscent of the coherent production mechanism of the axion. Models with low compactification scales (less than about 10^9 GeV) possess radions which are stable on cosmological time scales, for which there are strong overclosure constraints. When inflation ends, energy density is stored in radion oscillations which can easily dominate the energy density of the universe at late times. This strong moduli problem requires that, for models of conventional inflation, the inflation scale must be below the compactification scale $r \gtrsim 1$.

An alternative is the possibility that the radion could account for the dark matter required by cosmological measurements. Radion dark matter is probably challenging to detect directly, though evidence might be available from the gamma-ray spectrum (through the eventual, rare radion decay) or from isocurvature perturbations in the CMB.

We have examined the possibility that the radion could play the role of either the curvaton or modulon, and find that generally it is not suitable. It typically does not have large enough density fluctuations to be either a curvaton (unless it is already the inflaton) or a modulon. Finally, we have explored the possibility that the radion could decay after nucleosynthesis, and examined the stringent bounds which result. These bounds require both a much smaller abundance from coherent production and require a low reheat temperature to avoid thermal production mechanisms.

8.1 TeV Extra Dimensions

Models with TeV^{-1} extra dimensions have attracted much attention lately. Model builders in particle theory have made use of TeV^{-1} extra dimensions to suggest new mechanisms for breaking symmetries, naturally generating small numbers, and addressing the hierarchy problem [2, 3, 4] using orbifold boundary conditions. The natural scale for which these mechanisms to operate is the natural scale of physics beyond the SM. Thus, compactification

should be at the TeV scale⁷. Thus far, little has been said⁸ about the cosmology of these models, aside from the possibility of Kaluza–Klein dark matter, which is only an option in the special case of universal extra dimensions.

In the present work, we have derived additional and stronger constraints from the radion physics. We showed that TeV^{-1} extra dimensions generally have an overclosure problem, because the radion is effectively stable and easily dominates the energy density of the universe at late times. The radion energy density can be sufficiently damped provided the inflation scale M_I is at a TeV. The radion being stable, reheating would have to come from a different field which would imprint its fluctuations into the radiation.

To summarize, in models with TeV^{-1} extra dimensions:

- The radion is effectively stable.
- To avoid overclosure of the universe by radions, either the inflationary model must be rethought, or the inflation scale must be below the TeV scale *i.e.*, $r \gtrsim 1$. In this case, V_ϕ is subdominant during slow-roll and ϕ cannot be the inflaton. There is a narrow window of parameter space in which the radion can account for the observed abundance of dark matter.
- Inflation model building is challenging: A reheat temperature of order 100 GeV is preferred for baryogenesis and also for dark matter (unless the radion is dark matter—the LSP or LKP dark matter candidates actually require $T_{RH} \sim 100$ GeV). For $T_{RH} \sim \text{TeV} \sim M_I$, this necessitates a very efficient reheating, and almost certainly preheating.

In the last years, particle physics model building has moved into theories with TeV extra dimensions. We have explored the cosmological challenges associated with these models, and by seeing where constraints arise, define the properties that a solution of these challenges should have. One way out of these constraints points towards non-toroidal compactifications, like, for instance, warped geometries [19, 20, 21] or internal compact hyperbolic spaces for which $m_r \sim R^{-1}$ [39]. The future of extra-dimensional cosmology is both challenging and exciting.

Acknowledgments

G.S. thanks C. Armendáriz-Picón for useful discussions and T.T. is grateful for conversations with E. Pontón and to the SLAC theory group for hospitality while part of this work was

⁷We note that because the radion potential is protected by general covariance from large quantum corrections, the choice of M_c is stable with respect to M_* , the higher dimensional Planck scale, and thus is technically natural. However, the usual hierarchy problem, M_{EW} compared to M_* , remains somewhat mysterious in TeV extra dimensions (however, see [3, 4]).

⁸An observation related to the cosmology of TeV extra dimensions was made in [38] where it was found that there is a domain-wall problem in models with bulk $SU(N)$ gauge fields and fundamental brane matter if the reheat temperature is larger than M_c .

undertaken. This work is supported in part by the US Department of Energy, High Energy Physics Division, under contract W-31-109-Eng-38, by the David and Lucile Packard Foundation, and by NASA grant NAG5-10842.

References

- [1] I. Antoniadis, Phys. Lett. B **246**, 377 (1990).
- [2] N. Arkani-Hamed and M. Schmaltz, Phys. Rev. D **61**, 033005 (2000) [arXiv:hep-ph/9903417]; G. R. Dvali and M. A. Shifman, Phys. Lett. B **475**, 295 (2000) [arXiv:hep-ph/0001072]; D. E. Kaplan and T. M. Tait, JHEP **0111**, 051 (2001) [arXiv:hep-ph/0110126].
- [3] N. Arkani-Hamed, H. C. Cheng, B. A. Dobrescu and L. J. Hall, Phys. Rev. D **62**, 096006 (2000) [arXiv:hep-ph/0006238]. B. A. Dobrescu and E. Poppitz, Phys. Rev. Lett. **87**, 031801 (2001) [arXiv:hep-ph/0102010]; T. Appelquist, B. A. Dobrescu, E. Ponton and H. U. Yee, Phys. Rev. Lett. **87**, 181802 (2001) [arXiv:hep-ph/0107056]; H. J. He, C. T. Hill and T. M. Tait, Phys. Rev. D **65**, 055006 (2002) [arXiv:hep-ph/0108041]; T. Appelquist, B. A. Dobrescu, E. Ponton and H. U. Yee, Phys. Rev. D **65**, 105019 (2002) [arXiv:hep-ph/0201131].
- [4] I. Antoniadis, S. Dimopoulos, A. Pomarol and M. Quiros, Nucl. Phys. B **544**, 503 (1999) [arXiv:hep-ph/9810410]; A. Delgado, A. Pomarol and M. Quiros, Phys. Rev. D **60**, 095008 (1999) [arXiv:hep-ph/9812489]; D. E. Kaplan and T. M. Tait, JHEP **0006**, 020 (2000) [arXiv:hep-ph/0004200]; N. Arkani-Hamed, L. J. Hall, Y. Nomura, D. R. Smith and N. Weiner, Nucl. Phys. B **605**, 81 (2001) [arXiv:hep-ph/0102090]; C. Csaki, C. Grojean and H. Murayama, Phys. Rev. D **67**, 085012 (2003) [arXiv:hep-ph/0210133]; G. Burdman and Y. Nomura, Nucl. Phys. B **656**, 3 (2003) [arXiv:hep-ph/0210257].
- [5] G. Servant and T. M. Tait, Nucl. Phys. B **650**, 391 (2003) [arXiv:hep-ph/0206071]; H. C. Cheng, J. L. Feng and K. T. Matchev, Phys. Rev. Lett. **89**, 211301 (2002) [arXiv:hep-ph/0207125]; G. Servant and T. M. Tait, New J. Phys. **4**, 99 (2002) [arXiv:hep-ph/0209262]; D. Hooper and G. D. Kribs, Phys. Rev. D **67**, 055003 (2003) [arXiv:hep-ph/0208261]; G. Bertone, G. Servant and G. Sigl, arXiv:hep-ph/0211342; E. W. Kolb and R. Slansky, Phys. Lett. **B135**, 378 (1984).
- [6] T. Appelquist, H. C. Cheng and B. A. Dobrescu, Phys. Rev. D **64**, 035002 (2001) [arXiv:hep-ph/0012100].
- [7] M. Fairbairn, L. Lopez-Honorez and M. H. Tytgat, Phys. Rev. D **67**, 101302 (2003) [arXiv:hep-ph/0302160]. C. Armendariz-Picon and V. Duvvuri, arXiv:hep-th/0305237. A. Mazumdar and A. Perez-Lorenzana, Phys. Lett. B **508**, 340 (2001) [arXiv:hep-ph/0102174].
- [8] R. S. Chivukula, D. A. Dicus, H. J. He and S. Nandi, Phys. Lett. B **562**, 109 (2003) [arXiv:hep-ph/0302263].
- [9] N. Arkani-Hamed, S. Dimopoulos, N. Kaloper and J. March-Russell, Nucl. Phys. B **567**, 189 (2000) [arXiv:hep-ph/9903224].
- [10] C. L. Bennett *et al.*, Astrophys. J. **464**, L1 (1996) [arXiv:astro-ph/9601067].
- [11] D. H. Lyth and D. Wands, Phys. Lett. B **524**, 5 (2002) [arXiv:hep-ph/0110002]; D. H. Lyth, C. Ungarelli and D. Wands, Phys. Rev. D **67**, 023503 (2003) [arXiv:astro-ph/0208055].

- [12] G. Dvali, A. Gruzinov and M. Zaldarriaga, arXiv:astro-ph/0303591; L. Kofman, arXiv:astro-ph/0303614.
- [13] T. Appelquist and A. Chodos, Phys. Rev. D **28**, 772 (1983).
- [14] Z. Chacko and E. Perazzi, arXiv:hep-ph/0210254.
- [15] I. Antoniadis, K. Benakli, A. Laugier and T. Maillard, arXiv:hep-ph/0211409.
- [16] L. Perivolaropoulos and C. Sourdis, Phys. Rev. D **66**, 084018 (2002) [arXiv:hep-ph/0204155].
- [17] K. m. Cheung and G. Landsberg, Phys. Rev. D **65**, 076003 (2002) [arXiv:hep-ph/0110346];
T. Appelquist and H. U. Yee, Phys. Rev. D **67**, 055002 (2003) [arXiv:hep-ph/0211023].
A. J. Buras, M. Spranger and A. Weiler, Nucl. Phys. B **660**, 225 (2003) [arXiv:hep-ph/0212143].
A. J. Buras, A. Poschenrieder, M. Spranger and A. Weiler, arXiv:hep-ph/0306158.
- [18] N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **429**, 263 (1998) [arXiv:hep-ph/9803315]; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. R. Dvali, Phys. Lett. B **436**, 257 (1998) [arXiv:hep-ph/9804398].
- [19] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999) [arXiv:hep-ph/9905221].
- [20] W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83**, 4922 (1999) [arXiv:hep-ph/9907447].
- [21] W. D. Goldberger and M. B. Wise, Phys. Lett. B **475**, 275 (2000) [arXiv:hep-ph/9911457];
C. Csaki, M. L. Graesser and G. D. Kribs, Phys. Rev. D **63**, 065002 (2001) [arXiv:hep-th/0008151].
- [22] C. Csaki, M. Graesser, L. Randall and J. Terning, Phys. Rev. D **62**, 045015 (2000) [arXiv:hep-ph/9911406].
- [23] G. F. Giudice, R. Rattazzi and J. D. Wells, Nucl. Phys. B **595**, 250 (2001) [arXiv:hep-ph/0002178]; J. L. Hewett and T. G. Rizzo, arXiv:hep-ph/0202155; D. Dominici, B. Grzadkowski, J. F. Gunion and M. Toharia, arXiv:hep-ph/0206192.
- [24] L. Kofman, A. D. Linde and A. A. Starobinsky, Phys. Rev. D **56**, 3258 (1997) [arXiv:hep-ph/9704452].
- [25] N. Arkani-Hamed, H. C. Cheng, P. Creminelli and L. Randall, arXiv:hep-th/0301218.
- [26] L. Perivolaropoulos, arXiv:hep-ph/0301237.
- [27] C. Gordon and A. Lewis, arXiv:astro-ph/0212248.
- [28] H. Collins and R. Holman, Phys. Rev. D **67**, 105004 (2003) [arXiv:hep-ph/0210110].
- [29] J. R. Ellis, G. B. Gelmini, J. L. Lopez, D. V. Nanopoulos and S. Sarkar, Nucl. Phys. B **373** (1992) 399; R. H. Cyburt, J. R. Ellis, B. D. Fields and K. A. Olive, arXiv:astro-ph/0211258.
- [30] S. A. Abel and S. Sarkar, Phys. Lett. B **342**, 40 (1995) [arXiv:hep-ph/9409350].
- [31] J. L. Feng, A. Rajaraman and F. Takayama, arXiv:hep-ph/0302215.
- [32] G. F. Giudice, I. Tkachev and A. Riotto, JHEP **9908**, 009 (1999) [arXiv:hep-ph/9907510].
- [33] G. D. Kribs and I. Z. Rothstein, Phys. Rev. D **55**, 4435 (1997) [Erratum-ibid. D **56**, 1822 (1997)] [arXiv:hep-ph/9610468].

- [34] E. Ponton and E. Poppitz, JHEP **0106**, 019 (2001) [arXiv:hep-ph/0105021].
- [35] R. Sundrum, Phys. Rev. D **59**, 085010 (1999) [arXiv:hep-ph/9807348].
- [36] S. M. Carroll, J. Geddes, M. B. Hoffman and R. M. Wald, Phys. Rev. D **66**, 024036 (2002) [arXiv:hep-th/0110149].
- [37] N. Arkani-Hamed, L. J. Hall, D. R. Smith and N. Weiner, Phys. Rev. D **62**, 105002 (2000) [arXiv:hep-ph/9912453].
- [38] C. P. Korthals Altes and M. Laine, Phys. Lett. B **511**, 269 (2001) [arXiv:hep-ph/0104031].
K. Farakos, P. de Forcrand, C. P. Korthals Altes, M. Laine and M. Vettorazzo, Nucl. Phys. B **655**, 170 (2003) [arXiv:hep-ph/0207343].
- [39] N. Kaloper, J. March-Russell, G. D. Starkman and M. Trodden, Phys. Rev. Lett. **85**, 928 (2000) [arXiv:hep-ph/0002001].